

# RHEUMATOLOGY REFERRAL FORM

www.albertsons.com/specialtycare • Phone: 877.466.8028 • Fax: 877.466.8040



Patient Information

Patient Name: \_\_\_\_\_ DOB: \_\_\_\_\_ Sex: ☐ M ☐ F  
 Phone: \_\_\_\_\_ Cell Phone: \_\_\_\_\_ Email Address: \_\_\_\_\_  
 Address: \_\_\_\_\_ City: \_\_\_\_\_ State: \_\_\_\_\_ Zip: \_\_\_\_\_  
 ICD-10 Diagnosis Code: \_\_\_\_\_ Diagnosis: \_\_\_\_\_  
 Allergies (please note reaction): \_\_\_\_\_ ☐ Latex  
 Current Medications: (list here or attach a medication list): \_\_\_\_\_  
 Comorbidities: (list here or attach a list): \_\_\_\_\_

## INSURANCE INFORMATION - FAX COPY OF PATIENT'S INSURANCE CARD - BOTH SIDES

Prescription Information

MEDICATION	STRENGTH	DIRECTIONS	QUANTITY	REFILLS
<input type="checkbox"/> Actemra (tocilizumab)	162mg/0.9mL Prefilled Syringe	<input type="checkbox"/> Inject 162mg subcutaneously every other week. <input type="checkbox"/> Inject 162mg subcutaneously once a week.	28-day supply	
<input type="checkbox"/> Cimzia (certolizumab)	<input type="checkbox"/> Starter Kit 200mg/mL Prefilled Syringe <input type="checkbox"/> 200mg/mL Prefilled Syringe	<input type="checkbox"/> Loading Dose: Inject 400mg (2 syringes) subcutaneously at weeks 0, 2 and 4. <input type="checkbox"/> Maintenance Dose: Inject 400mg (2 syringes) subcutaneously every 4 weeks. <input type="checkbox"/> Maintenance Dose: Inject 200mg (1 syringe) subcutaneously every 2 weeks.	1 kit (6 syringes) 28-day supply	
<input type="checkbox"/> Cosentyx (secukinumab)	<input type="checkbox"/> 150mg Sensoready Pen <input type="checkbox"/> 150mg Prefilled Syringe	<input type="checkbox"/> Loading Dose: Inject 150mg subcutaneously once weekly at weeks 0, 1, 2, 3, and 4. <input type="checkbox"/> Loading Dose: Inject 300mg (2 injections of 150mg) subcutaneously once weekly at weeks 0, 1, 2, 3, and 4. <input type="checkbox"/> Maintenance Dose: Inject 150mg subcutaneously every 4 weeks. <input type="checkbox"/> Maintenance Dose: Inject 300mg (2 injections of 150mg) subcutaneously every 4 weeks.	5 doses 28-day supply	
<input type="checkbox"/> Enbrel (etanercept)	<input type="checkbox"/> 50mg/mL Sureclick Auto-injector <input type="checkbox"/> 50mg/mL Prefilled Syringe <input type="checkbox"/> 25mg/0.5mL Prefilled Syringe <input type="checkbox"/> 25mg Vial	<input type="checkbox"/> Inject 50mg subcutaneously once a week. <input type="checkbox"/> Inject 25mg subcutaneously twice a week.	28-day supply 28-day supply	
<input type="checkbox"/> Humira (adalimumab)	10kg to < 15kg <input type="checkbox"/> 10mg/0.1 mL Citrate-free Prefilled Syringe <input type="checkbox"/> 10mg/0.2 mL Prefilled Syringe 15kg to < 30kg <input type="checkbox"/> 20mg/0.2 mL Citrate-free Prefilled Syringe <input type="checkbox"/> 20mg/0.4 mL Prefilled Syringe ≥ 30kg <input type="checkbox"/> 40mg/0.4 mL Citrate-free Pen <input type="checkbox"/> 40mg/0.8 mL Pen <input type="checkbox"/> 40mg/0.4 mL Citrate-free Prefilled Syringe <input type="checkbox"/> 40mg/0.8 mL Prefilled Syringe	<input type="checkbox"/> Inject 10mg subcutaneously every OTHER week. <input type="checkbox"/> Inject 20mg subcutaneously every OTHER week. <input type="checkbox"/> Inject 40mg subcutaneously every OTHER week. <input type="checkbox"/> Inject 40mg subcutaneously every week.	28-day supply	
<input type="checkbox"/> Kineret (anakinra)	100mg/0.67mL Prefilled Syringe	Inject 100mg subcutaneously every 24 hours.	28-day supply	
<input type="checkbox"/> Orencia (abatacept)	<input type="checkbox"/> 250mg Vial (IV use only) <input type="checkbox"/> 125mg/mL Prefilled Syringe <input type="checkbox"/> 125mg/mL Auto-injector <input type="checkbox"/> 87.5/0.7mL Prefilled Syringe <input type="checkbox"/> 50mg/0.4mL Prefilled Syringe	<input type="checkbox"/> Loading Dose: Inject _____mg via intravenous infusion at 0, 2 and 4 weeks. <input type="checkbox"/> Inject _____mg via intravenous infusion every 4 weeks. <input type="checkbox"/> Adults and children ≥ 50 kg: Inject 125mg subcutaneously once weekly. <input type="checkbox"/> Children ≥ 25 to < 50 kg: Inject 87.5mg subcutaneously once weekly. <input type="checkbox"/> Children 10 to < 25 kg: Inject 50mg subcutaneously once weekly.	28-day supply 28-day supply 28-day supply	

Prescription information continued on next page

MEDICATION	STRENGTH	DIRECTIONS	QUANTITY	REFILLS
<input type="checkbox"/> Otrexup (methotrexate)	<input type="checkbox"/> 7.5mg/0.4mL Auto-injector <input type="checkbox"/> 10mg/0.4mL Auto-injector <input type="checkbox"/> 12.5mg/0.4mL Auto-injector <input type="checkbox"/> 15mg/0.4mL Auto-injector <input type="checkbox"/> 17.5mg/0.4mL Auto-injector <input type="checkbox"/> 20mg/0.4mL Auto-injector <input type="checkbox"/> 22.5mg/0.4mL Auto-injector <input type="checkbox"/> 25mg/0.4mL Auto-injector	Inject one auto-injector subcutaneously once weekly.	28-day supply	
<input type="checkbox"/> Rasuvo (methotrexate)	<input type="checkbox"/> 7.5mg/0.15 Auto-injector <input type="checkbox"/> 10mg/0.2mL Auto-injector <input type="checkbox"/> 12.5mg/0.25mL Auto-injector <input type="checkbox"/> 15mg/0.3mL Auto-injector <input type="checkbox"/> 17.5mg/0.35mL Auto-injector <input type="checkbox"/> 20mg/0.4mL Auto-injector <input type="checkbox"/> 22.5 mg/0.45mL Auto-injector <input type="checkbox"/> 25mg/0.5mL Auto-injector <input type="checkbox"/> 27.5mg/0.55mL Auto-injector <input type="checkbox"/> 30mg/0.6mL Auto-injector	Inject one auto-injector subcutaneously once weekly.	28-day supply	
<input type="checkbox"/> Remicade (infliximab) <input type="checkbox"/> Inflectra (infliximab-dyyb)	100mg Vial	<input type="checkbox"/> Loading Dose: Administer _____mg (at at _____mg/kg) intravenously at 0, 2 and 6 weeks. <input type="checkbox"/> Maintenance Dose: Administer _____mg (at _____mg/kg) intravenously every _____weeks.	42-day supply 28-day supply	
<input type="checkbox"/> Simponi (golimumab)	<input type="checkbox"/> 50mg/0.5mL SmartJect Auto-injector <input type="checkbox"/> 50mg/0.5mL Prefilled Syringe	Inject 50mg subcutaneously once monthly.	28-day supply	
<input type="checkbox"/> Taltz (ixekizumab)	<input type="checkbox"/> 80mg/mL Auto-injector <input type="checkbox"/> 80mg/mL Prefilled Syringe	<input type="checkbox"/> Inject 160mg subcutaneously at weeks 2, 4, 6, 8, 10 and 12; then inject 80mg subcutaneously every 4 weeks. <input type="checkbox"/> Inject 160mg subcutaneously once, followed by 80mg subcutaneously every 4 weeks. <input type="checkbox"/> Inject 80mg subcutaneously every 4 weeks.	<input type="checkbox"/> 28-day supply <input type="checkbox"/> 84-day supply	
<input type="checkbox"/> Xeljanz (tofacitinib)	<input type="checkbox"/> 5mg Tablet	<input type="checkbox"/> Take 1 tablet by mouth twice daily. <input type="checkbox"/> Take 1 tablet by mouth once a day (renal/hepatic impairment).	60 30	
<input type="checkbox"/> Xeljanz XR (tofacitinib)	<input type="checkbox"/> 11mg XR Tablet	Take 1 tablet by mouth once daily.	30	
Other Medication Name:				

**Treatment History:** ☐ **New to Therapy** ☐ **Continuation of Therapy**

Hepatitis B Screening Results: ☐ HBsAg: \_\_\_\_\_ ☐ Anti-HBs: \_\_\_\_\_ ☐ Anti-HBc: \_\_\_\_\_  
 If applicable, has treatment been initiated? ☐ Yes ☐ No  
 Tuberculosis Assessment Date: ☐ Negative ☐ Active TB ☐ Latent TB ☐ History of active or latent TB  
 If history of active or latent TB: \_\_\_\_\_ Adequate treatment is confirmed: ☐ Yes ☐ No  
 History of Irritable Bowel Disease: ☐ Yes ☐ No

Prescriber Name: \_\_\_\_\_  
 State License #: \_\_\_\_\_ DEA #: \_\_\_\_\_ NPI: \_\_\_\_\_  
 Additional Contact Person Name: \_\_\_\_\_  
 Group or Hospital: \_\_\_\_\_ Phone: \_\_\_\_\_  
 Fax: \_\_\_\_\_ Email Address: \_\_\_\_\_  
 Address: \_\_\_\_\_ City: \_\_\_\_\_ State: \_\_\_\_\_ Zip: \_\_\_\_\_  
 Prescriber Signature: \_\_\_\_\_  

Product Substitution Permitted
Dispensed as Written
Date

☐ Ship to Patient ☐ Ship to Prescriber/Clinic ☐ Pick up at Albertsons Companies Pharmacy  
 Date Medication Needed: \_\_\_\_\_