## ONCOLOGY REFERRAL FORM

www.albertsons.com/specialtycare · Phone: 877.466.8028 · Fax: 877.466.8040









VONS.

ACME.



PAVILIONS CARRS () Randalls Tom Thumb

	_
	_
44	
_	.≃
	- 1
-	_
a	- cc
-	_
	_
10	_
•	
-	4
	_
	_

Prescription

Patient Name:			_ DOB:			Sex: M	F
Phone:				Email Address:			
Address:		City:			State:	Zip:	
ICD-10 Diagnosis Code:		Diagnosis:					
Allergies (please note reaction):							Latex
Current Medications: (list here or attach a	medication list):						
Comorbidities: (list here or attach a list):							

## **INSURANCE INFORMATION - FAX COPY OF PATIENT'S INSURANCE CARD - BOTH SIDES**

MEDICATION	STRENGTH	DIRECTIONS	QUANTITY	REFILLS
Bosulif (bosutinib)	☐ 100mg Tab	☐ Take 500mg by mouth once daily with food.	20 day ayaaly	
	☐ 500mg Tab	☐ Takemg by mouth once daily with food.	30-day supply	
	☐ 75mg Cap			
☐ Ibrance (palbociclib)	☐ 100mg Cap	Take 1 capsule by mouth once daily with food for 21 days, followed by 7 days off for a 28 day cycle.	21 caps	
	☐ 125mg Cap			
☐ WITH letrozole	2.5mg Tab	Take 1 tablet by mouth once daily.	30-day supply	
☐ WITH anastrozole	1mg Tab	Take 1 tablet by mouth once daily.	30-day supply	
☐ WITH exemestane	25mg Tab	Take 1 tablet by mouth once daily after a meal.	30-day supply	
☐ WITH fulvestrant	500mg Sol	☐ Inject 500mg intramuscularly on days 1, 15, 29 then monthly thereafter.	30-day supply	
☐ Inlyta (axitinib)	5mg Tab	☐ Take 5mg by mouth twice daily, approximately 12 hours apart.	20 day ayarda	
	1mg Tab	Takemg by mouth twice daily, approximately 12 hours apart.	30-day supply	
☐ Keytruda (pembrolizumab)	50mg lyophilized powder in SDV for reconstitution	Administer 200mg via intravenous infusion every 3 weeks over 30 minutes.	21-day supply	
	100 mg/4ml solution in single-dose vial	Pediatrics: At 2mg/kg (max. 200mg), administermg via intravenous infusion every 3 weeks over 30 minutes.V		
☐ Kisqali (ribociclib)	200 mg Tala	Take 600mg (3 tablets) by mouth once daily for 21 days followed by 7 days off treatment.	28-day supply	
	200mg Tab	Takemg by mouth once daily for 21 days followed by 7 days off treatment	28-day supply	
☐ WITH letrozole	2.5 mg Tab	☐ Take 1 tablet by mouth twice daily.	30-day supply	
Kisqali Femara Co-pack (ribociclib and letrozole)	200mg Tab/ 2.5mg Tab	Take Kisqali 600mg (3 tablets) by mouth once daily for 21 consecutive days followed by 7days off treatment and take Femara 2.5 mg by mouth once daily continuously for a 28-day cycle.	28-day supply	

Prescription information continued on next page

## It's as simple as caring.





















